

LONGITUDINAL CHANGES IN RESPONSE TO A CYCLE-RUN FIELD TEST OF YOUNG MALE NATIONAL “TALENT IDENTIFICATION” AND SENIOR ELITE TRIATHLON SQUADS

VÍCTOR DÍAZ,^{1,2} ANA B. PEINADO,² VERONICA E. VLECK,³ MARÍA ALVAREZ-SÁNCHEZ,¹ PEDRO J. BENITO,² FRANCISCO B. ALVES,³ FRANCISCO J. CALDERÓN,² AND AUGUSTO G. ZAPICO⁴

¹Institute of Veterinary Physiology, University of Zürich and Zürich Center for Integrative Human Physiology (ZIHP), Zürich, Switzerland; ²Department of Health and Human Performance, School of Physical Activity and Sport Sciences-INEF, Technical University of Madrid, Madrid, Spain; ³CIPER, Faculty of Human Kinetics, Technical University of Lisbon, Cruz Quebrada-Dafundo, Portugal; and ⁴Faculty of Education, Complutense University of Madrid, Madrid, Spain

ABSTRACT

Díaz, V, Peinado, AB, Vleck, VE, Alvarez-Sánchez, M, Benito, PJ, Alves, FB, Calderón, FJ, and Zapico, AG. Longitudinal changes in response to a cycle-run field test of young male national “talent identification” and senior elite triathlon squads. *J Strength Cond Res* 26(8): 2209–2219, 2012—This study investigated the changes in cardiorespiratory response and running performance of 9 male “Talent Identification” (TID) and 6 male Senior Elite (SE) Spanish National Squad triathletes during a specific cycle-run (C-R) test. The TID and SE triathletes (initial age 15.2 ± 0.7 vs. 23.8 ± 5.6 years, $p = 0.03$; $\dot{V}O_2\text{max}$ 77.0 ± 5.6 vs. 77.8 ± 3.6 ml·kg⁻¹·min⁻¹, nonsignificant) underwent 3 tests through the competitive period and the preparatory period, respectively, of 2 consecutive seasons: test 1 was an incremental cycle test to determine the ventilatory threshold (Th_{vent}); test 2 (C-R) was 30-minute constant load cycling at the Th_{vent} power output followed by a 3-km time-trial run; and test 3 (isolated control run [R]) was an isolated 3-km time-trial control run, in randomized counter-balanced order. In both seasons, the time required to complete the C-R 3-km run was greater than for R in TID ($11:09 \pm 00:24$ vs. $10:45 \pm 00:16$ min:ss, $p < 0.01$ and $10:24 \pm 00:22$ vs. $10:04 \pm 00:14$, $p = 0.006$, for season 2005–2006 and 2006–2007, respectively) and SE ($10:15 \pm 00:19$ vs. $09:45 \pm 00:30$, $p < 0.001$ and $09:51 \pm 00:26$ vs. $09:46 \pm 00:06$, $p = 0.02$ for season 2005–2006 and 2006–2007, respectively). Compared with the first season, the completion of the time-trial run was faster in the second season (6.6%, $p < 0.01$

and 6.4%, $p < 0.01$, for C-R and R tests, respectively) only in TID. Changes in post cycling run performance were accompanied by changes in pacing strategy, but there were only slight or nonsignificant changes in the cardiorespiratory response. Thus, the negative effect of cycling on performance may persist, independently of the period, over 2 consecutive seasons in TID and SE triathletes; however, improvements over time suggests that monitoring running pacing strategy after cycling may be a useful tool to control performance and training adaptations in TID.

KEY WORDS transition, performance, maximum oxygen consumption, pacing

INTRODUCTION

The running component of male Elite draft-legal Olympic triathlon is becoming more important for overall race performance. Correlation coefficients of >0.8 were observed between time to complete the run section in seconds and finishing position in male International Triathlon Union (ITU) World Cup, European Championship, and World Championship competition, respectively, in 2002, 2005, and 2006 ([38]; V.E. Vleck and A. Bürgi, personal communication, 2011). Overall triathlon performance is not only affected by running performance but also by performance during the cycle section. The higher the athlete is placed at the end of the cycle section, the greater the importance to his final position (29). Overall, the cycle to run transition (T2) is recognized as a crucial moment during competition, and consequently, it has been widely studied from both physiological and biomechanical points of view (see the bibliography in [35]).

Specific-T2 responses have been investigated using both laboratory and field-based tests. Running after cycling has been shown to elicit increases in oxygen consumption ($\dot{V}O_2$) (22), ventilation (\dot{V}_E) (18,20), and heart rate (HR) (22) when

Address correspondence to Víctor Díaz, diaz@vetphys.uzh.ch.

26(8)/2209–2219

Journal of Strength and Conditioning Research

© 2012 National Strength and Conditioning Association

TABLE 1. Characteristics of the participants.*

	Senior elite (n = 6)		Talent identification (n = 9)	
	Season 2005–2006	Season 2006–2007	Season 2005–2006	Season 2006–2007
Age (y)	24 ± 5.6	24.8 ± 5.6	15.2 ± 0.7†	16.2 ± 0.7†
Mass (kg)	71.2 ± 8.7	71.9 ± 6.8	60.2 ± 6.8	61.9 ± 6.6
Height (cm)	180.0 ± 8.8	180.2 ± 8.6	173.6 ± 6.4	174.0 ± 7.3
% Fat	8.5 ± 0.6	8.3 ± 0.4	7.7 ± 0.6	7.9 ± 0.7
$\dot{V}O_2\text{max}$ (ml·kg ⁻¹ ·min ⁻¹)	77.8 ± 3.6	77.4 ± 4.6	77.0 ± 5.0	76.8 ± 5.9
HR _{max} (b·min ⁻¹)	186 ± 3	184 ± 4	190 ± 2	188 ± 4
PO _{max} (W·kg ⁻¹)	5.7 ± 1.2	5.9 ± 0.8	5.1 ± 1.0†	5.3 ± 1.1†
Th _{vent} (% $\dot{V}O_2\text{max}$)	64.8 ± 3.1	66.1 ± 4.0	61.9 ± 5.2	63.1 ± 5.5
Th _{vent} (W·kg ⁻¹)	3.4 ± 0.8	3.6 ± 1.0	3.5 ± 1.1	3.6 ± 0.9

*HR_{max} = maximal heart rate; $\dot{V}O_2\text{max}$ = cycling maximum oxygen uptake; HR_{max} = maximum heart rate; PO_{max} = maximum power output; Th_{vent} = ventilatory threshold.

†Differences (p < 0.05) with Senior Elite triathletes.

compared with an isolated control run. The extent to which such increases occur, for both competition and controlled conditions, appears to be influenced by the athlete’s ability (25,27,28) and training status (5,12,27,28). It has been suggested that the assessment of “specific-T2 adaptation” over time, via the examination of performance related indices (such as mean running speed or the metabolic cost of running) within the post cycling-running component of

a “simulated T2” test, may prove beneficial to the battery of tests usually performed by young (potential Elite) triathletes (25). Moreover, in a later study Vleck et al. (36) suggested that cardiorespiratory variables may be a more sensitive gauge of T2-specific adaptation than blood lactate concentration is. They also highlighted the need for a T2-related test that is sensitive enough to detect changes in specific-T2 adaptive response over time. In addition to the potential

TABLE 2. Cardiorespiratory response and running performance (mean ± SD) in talent identification triathletes (n = 9) during the C-R and R tests in season 2005–2006 and 2006–2007.*

	Season 2005–2006		Season 2006–2007	
	C-R	R	C-R	R
$\dot{V}O_2$ (mL·min ⁻¹)	4,222 ± 520	4,350 ± 707	4,549 ± 623†	4,088 ± 784
$\dot{V}O_2$ (% cycling $\dot{V}O_2\text{max}$)	89.9 ± 4.3	90.1 ± 3.9	92.1 ± 3.1†	87.5 ± 6.1
$\dot{V}CO_2$ (mL·min ⁻¹)	3,615 ± 298	3,639 ± 604	3,855 ± 468	3,563 ± 518
\dot{V}_E (L·min ⁻¹)	132.5 ± 16.8	127.2 ± 24.3‡	139.5 ± 13.8	124.7 ± 23.8‡
$\dot{V}_E/\dot{V}O_2$	35.0 ± 4.0	32.3 ± 2.8‡	33.4 ± 1.9†	30.6 ± 2.0†‡
$\dot{V}_E/\dot{V}CO_2$	35.6 ± 3.7	34.0 ± 3.3‡	35.2 ± 1.4	30.7 ± 3.6 †‡
f (resp·min ⁻¹)	57 ± 10	55 ± 11	60 ± 4	58 ± 9
HR (b·min ⁻¹)	186 ± 8	181 ± 12‡	184 ± 8	175 ± 10†‡
RER	0.87 ± 0.04	0.85 ± 0.04‡	0.84 ± 0.02	0.90 ± 0.07†‡
RPE	9.5 ± 0.5	9.0 ± 0.5	9.4 ± 0.5	9.0 ± 0.5
Cadence (rpm)	91 ± 2		90 ± 2	
Speed (km·h ⁻¹)	16.3 ± 1.1 (3:43)§	16.7 ± 0.7‡ (3:35)	17.3 ± 0.7† (3:28)	17.8 ± 0.4†‡ (3:21)
Time to complete run (mm:ss)	11:09 ± 00:24	10:45 ± 00:16‡	10:24 ± 00:22†	10:04 ± 00:14†‡

*HR = heart rate; RER = respiratory exchange ratio; RPE = rating of perceived exertion; f = breathing frequency; $\dot{V}O_2$ = oxygen uptake; $\dot{V}CO_2$ = carbon dioxide production; \dot{V}_E = minute ventilation.

†Differences (p < 0.05) with season 2005–2006 within the test.

‡Differences (p < 0.05) with C-R within the season.

§Average speed or pace is shown between parentheses as minutes:seconds:per kilometer.




Figure 1. Change in time to complete the 3-km time-trial run during season 2005–2006 and 2006–2007 in Talent Identification (left panels, $n = 9$) and Senior Elite (right panels, $n = 6$) triathletes. Data are normalized to the isolated running (R) test. The dashed line represents the mean of the group. *Differences ($p < 0.05$) between tests.

ability of such test to distinguish between groups of athletes with different performance levels, this sensitivity would be important in the context of development of young triathletes.

Hitherto, no investigations regarding the longitudinal changes in performance and cardiorespiratory response to running after cycling have been published. The extent of sensitivity to changes in the T2 adaptation over time between tests is also unknown, and this may prove to be a useful tool in monitoring young athletes who are at the “talent identification” (TID) stage of their career. Therefore, the aim of this study was to assess the extent to which the cardiorespiratory response and running performance change over the course of 2 consecutive race seasons using a simulated T2 test in 2 different ability and maturation groups (34) of triathletes, that is, the best Spanish under-16 year-old boys and men within the Spanish Senior Elite (SE) National Squad.

METHODS

Experimental Approach to the Problem

We evaluated, in 2 different groups (see below), the specific-T2 cardiorespiratory response and running performance at

the same period of 2 consecutive seasons (2005–2006 and 2006–2007). The experimental protocol involved a preliminary test to determine maximal oxygen uptake ($\dot{V}O_{2max}$) during cycling ergometry and the individual ventilatory threshold (Th_{vent}), followed by 2 tests in randomized counterbalanced order: (a) a field-based cycle-run (C-R) test and (b) an isolated control run (R), as described in detail below. The 2 different tests, C-R and R, were completed on an outdoor 400-m synthetic Tartan track. Running performance (time to complete the 3-km time trial) and cardiorespiratory response were measured during the C-R and R tests to evaluate both the changes over 2 consecutive seasons and the negative effect of cycling on subsequent running.

Subjects

Two groups of male triathletes participated in the study (Table 1). The first group was composed of 9 young triathletes who had been selected by the Spanish Triathlon Federation (FETRI) as being the best in the country. They had 18 ± 8 months of previous triathlon experience and had previously obtained a top 10 finish in the under-16 category National




Figure 2. Lap-by-lap analysis of the cardiorespiratory response of Talent Identification triathletes ($n = 9$) during the isolated running (R) test (left panels) and the cycle-run (C-R) test (right panels) in season 2005–2006 (close circles) and 2006–2007 (open circles). Data represent the mean \pm SEM. *Differences ($p < 0.05$) between seasons within the test. #Differences ($p < 0.05$) between tests within the season.

Triathlon Championships 1 week before the start of the study. All of them were members of the FETRI *National Plan for Talent Identification* and are hereafter referred to as the TID group. Testing took place during a specific training camp in which athletes were living together, organized by FETRI, with all athletes following the same training schedules and the same

diet. All the triathletes passed a medical examination in the week before the experiments.

A second group of 6 SE Spanish National Squad triathletes took part in the study. The SE triathletes had 8.1 ± 2 years of previous training background and participated regularly in international competitions. All were based at the High Performance Center in Madrid and followed a similar diet and training schedules during the period of the study. The total number of athletes in the Senior National Squad fluctuated over the 2-year experiment, but at no time did our SE group account for $<55\%$ or for $>65\%$ of the complete male Spanish National Senior Squad. The SE triathletes were regularly assessed by the National team physician and had passed a complete medical examination at the beginning of the season (i.e., October).

In agreement with the guidelines of the 1975 Declaration of Helsinki, the study was approved by the Local University Ethics Committee. The participants were given both verbal and written information regarding the possible risks and benefits of the study. Signed informed consent was obtained from the senior triathletes and from the parents or legal guardians of the TID group before the start of the study.

Procedures

The TID triathletes carried out all the tests in Soria (Spain) during the competitive period (July). International racing commitments precluded the SE group from being tested at the equivalent period of the season,

and they were therefore tested (February of each respective year) in Madrid (Spain) during the final weeks of the preparatory phase. Both cities are at approximately 700-m altitude, the wind was calm, and the environmental conditions were both stable and similar in both seasons (Soria: $22 \pm 2.4^{\circ}\text{C}$, $58 \pm 3.8\%$, 702 ± 12 mm Hg during season 2005–2006 vs. $24 \pm$

TABLE 3. Cardiorespiratory response and running performance (mean \pm SD) in Senior Elite triathletes ($n = 6$) during the C-R and R tests in seasons 2005–2006 and 2006–2007.*

	Season 2005–2006		Season 2006–2007	
	C-R	R	C-R	R
$\dot{V}O_2$ (mL·min ⁻¹)	4,730 \pm 340	4,691 \pm 914	4,730 \pm 307	4,750 \pm 333
$\dot{V}O_2$ (% cycling $\dot{V}O_{2max}$)	87.0 \pm 3.6	84.5 \pm 5.6	85.7 \pm 4.5	88.2 \pm 4.0
$\dot{V}CO_2$ (mL·min ⁻¹)	4,119 \pm 258	5,145 \pm 319	5,186 \pm 221 [†]	4,055 \pm 203 [†]
\dot{V}_E (L·min ⁻¹)	139.8 \pm 11.0	151.3 \pm 18.7	155.6 \pm 16.4	137.6 \pm 6.8
$\dot{V}_E/\dot{V}O_2$	33.18 \pm 1.34	37.59 \pm 7.53	38.12 \pm 4.28 [†]	32.88 \pm 2.43
$\dot{V}_E/\dot{V}CO_2$	32.97 \pm 1.15	28.57 \pm 1.93	29.24 \pm 2.76 [†]	33.26 \pm 2.44 [†]
f (resp·min ⁻¹)	55 \pm 3	56 \pm 7	56 \pm 5	54 \pm 4
HR (b·min ⁻¹)	183 \pm 7	171 \pm 8 [‡]	175 \pm 9	178 \pm 5 [‡]
RER	0.88 \pm 0.03	1.03 \pm 0.08	1.02 \pm 0.05 [†]	0.87 \pm 0.01 [†]
RPE	9.2 \pm 0.4	9.0 \pm 0.5	9.3 \pm 0.5	9.0 \pm 0.6
Cadence (rpm)	94 \pm 3		95 \pm 2	
Speed (km·h ⁻¹)	17.5 \pm 0.5 (3:25) [§]	18.5 \pm 1.2 [‡] (3:15)	18.2 \pm 0.8 (3:17)	18.4 \pm 0.2 [‡] (3:15)
Time to complete run (mm:ss)	10:15 \pm 00:19	09:45 \pm 00:30 [‡]	09:51 \pm 00:26	09:46 \pm 00:06 [‡]

* \dot{V}_T = tidal volume; f = breathing frequency; HR = heart rate; RER = respiratory exchange ratio; RPE = rating of perceived exertion; $\dot{V}O_2$ = oxygen uptake; $\dot{V}CO_2$ = carbon dioxide production; \dot{V}_E = minute ventilation; f = breathing frequency; HR = heart rate; RER = respiratory exchange ratio; RPE = rating of perceived exertion.

[†]Differences ($p < 0.05$) with season 2005–2006 within the test.

[‡]Differences ($p < 0.05$) with C-R within the season.

[§]Average speed or pace is shown between parentheses as minutes:seconds-per kilometer.

2.1°C, 55 \pm 5.3% and 705 \pm 15 mm Hg during season 2006–2007. Madrid: 21.0 \pm 3.2°C, 47 \pm 2.1%, 699 \pm 9 mm Hg during season 2005–2006 vs. 21.4 \pm 3.0°C, 43 \pm 3.4%, 701 \pm 14 mm Hg during season 2006–2007). No significant intralocation or inter location differences in environmental conditions were noted at any point of the study.

All the tests were performed at the same time of the day, interspersed by 1 day of passive or active (swim only) rest. The subjects were constantly encouraged to give their maximum performance, but they were not given any time or HR data during the tests.

Determination of Cycling $\dot{V}O_{2max}$ and the Ventilatory Threshold

All the athletes performed an incremental test on a cycloergometer (Cardgirus[®] cycloergometer, G&G Innovación S.A., Madrid, Spain)—for which frame specifications were set by the athlete during his first test occasion to mimic his normal cycle position and maintained during the study. The test started with a 5-minute warm-up period at 75 W·min⁻¹. Load was then increased by 25 W·min⁻¹ until voluntary exhaustion. The athletes were asked to maintain a cadence from 70 to 90 rpm throughout the test. All the tests fulfilled at least 2 of the following criteria (31): A respiratory exchange ratio (RER) >1.10, a plateau in $\dot{V}O_2$ (corresponding to a variation of <100 ml·min⁻¹) despite an increase in exercise intensity and a peak HR >220 – age.

Breath-by-breath analysis of expired air was performed using a Jaeger Oxycon Mobile[®] (13,32) gas analyzer (Erich Jaeger, Viasys Healthcare, Hoechst, Germany). The analyzer was calibrated immediately before testing according to the manufacturer's instructions. The data were averaged every 30 seconds, and the highest value at the highest load achieved was termed $\dot{V}O_{2max}$.

The Th_{vent} was estimated on data averaged every 10 seconds, using the V-slope method (2) by 2 experienced researchers working independently. Because the coefficient of variation between said researchers' determinations of Th_{vent} was 1.8% and similar to that reported elsewhere (31), the intervention of a third researcher was not required.

The Cycle-Run Test

All the subjects carried out a 10-minute warm-up (involving 5-minute cycling at <100 W and 5-minute running at a very easy but freely chosen intensity) before the test. The C-R test comprised 30 minutes of steady-state cycling at the workload corresponding to Th_{vent} (16–18,20) determined in test 1 followed by a self-paced 3-km run (3,4). The subjects spent 34 \pm 8 seconds, on average, to complete the cycle ergometer-to-track transition and were instructed to complete the running section of the trial as fast as possible. All the subjects ingested 250 ml of cool water during the first 20 minutes of the cycling component of C-R. The portable gas analyzer (Oxycon Mobile, Erich Jaeger) was then attached to




Figure 3. Lap-by-lap analysis of the cardiorespiratory response of Senior Elite triathletes ($n = 6$) during the isolated running (R) test (left panels) and the cycle-run (C-R) test (right panels) in season 2005–2006 (close circles) and 2006–2007 (open circles). Data represent the mean \pm SEM. No differences between seasons within the test were found. #Differences ($p < 0.05$) between tests within the season.

variables were chosen because of the necessity to obtain >1–2 data points per 400-m lap. The HR was also recorded over 10-second intervals using an HR monitor (Polar 620i, Polar Electro, Kempele, Finland). All the variables, including speed (distance per time), were then expressed as mean \pm SD for consecutive laps of the run track.

Isolated Running Test

After a 10-minute warm-up at a “very easy” self-selected pace, the athletes were asked to perform a 3-km run time trial (3,4) as fast as possible. During the R test, the cardiorespiratory responses of the athletes were measured as in the C-R test by means of the portable gas analyzer. The subjects’ rating of perceived exertion (RPE) was recorded using the modified 10-point Borg scale (9) immediately after the completion of both R and C-R test.

Statistical Analyses

Differences in $\dot{V}O_2$, $\dot{V}CO_2$, \dot{V}_E , $\dot{V}_E/\dot{V}O_2$, $\dot{V}_E/\dot{V}CO_2$, RER, f , and HR between the running component of the C-R test and the R test were analyzed within the group via multivariate analysis with season (season 2005–2006 vs. season 2006–2007), lap and condition (C-R vs. R) as independent factors. When significant differences were detected, pairwise comparisons with Bonferroni Post hoc adjustment were performed. Given the sample size (9 young and 6 Seniors triathletes) the statistical power achieved was >80%.

Performance, defined as the time required to complete the 3-km time-trial run, was compared between tests within the season

(C-R vs. R) and between seasons within the test (season 2005–2006 vs. season 2006–2007), using paired Student t -tests. For all the analyses, significance was set at $p < 0.05$. All the calculations were made using SPSS v.18.0 software for

the triathletes’ back using a harness and $\dot{V}O_2$, CO_2 production ($\dot{V}CO_2$), \dot{V}_E , and respiratory frequency (f) were recorded and averaged over 10-second intervals. Ten seconds rather than the more customary 30-second averages for respiratory




Figure 4. Change in time to complete the 3-km time-trial run during the cycle-run (C-R) or isolated running (R) test in Talent Identification (left panels, $n = 9$) and Senior Elite (right panels, $n = 6$) triathletes. Data are normalized to season 2005–2006. The dashed line represents the mean of the group. *Differences ($p < 0.05$) between seasons.

Windows[®] (SPSS Worldwide Headquarters, Chicago, IL, USA).

RESULTS

Both groups of athletes exhibited a cycling $\dot{V}O_{2max}$ and peak power output values (Table 1) comparable with those reported previously for members of the French National Squad (3,19,25,26). Between January 2005 and October 2007, the average pace during race competitions (draft-legal) was recorded in a minimum of 4 and a maximum of 8 races per subject (mean \pm SD = 5.5 ± 1.1). All the competitions for SE triathletes were performed over 10-km running sections and included ITU or European Triathlon Union (ETU) events. For TID triathletes, competitions included 3- or 5-km (as the Spanish Championships 1 week previous to the tests) running segments. The average mean speed was 3:25 and 3:35 min:ss·km⁻¹ for SE and TID group, respectively.

Differences between Cycle-Run and Isolated Running Tests

In the TID triathletes, cycling had a negative effect on running performance in both seasons. The mean time taken to

complete the running component of C-R was longer than that needed to complete the R test, by 3.5% (or 24 seconds, $p < 0.01$) and 3.3% (or 20 seconds, $p < 0.01$) in the 2005–2006 and the 2006–2007 seasons, respectively (Table 2, Figure 1). The \dot{V}_E , the ratios of $\dot{V}_E/\dot{V}O_2$ and $\dot{V}_E/\dot{V}CO_2$, and the HR were also all higher ($p < 0.001$) in the C-R test than in the R test, in both seasons (Table 2, Figure 2).

Table 3 shows the results of the SE triathletes. The cardiorespiratory response of the SE during the running component of the C-R test did not differ from that exhibited within the isolated run, with the exception of the HR. During the C-R run, the HR was lower ($p < 0.01$) in the first season and higher ($p = 0.03$) in the second season of the study (Table 3, Figure 3). Within 400-m lap analyses also revealed significant differences in the HR between the C-R and the R test within the season, over the first 400 m (season 2005–2006) or 800 m (season 2006–2007) of the run time trial (Figure 3). Moreover, time to complete the time-trial run was also negatively affected (by 4.9% of total control run time, 30 seconds, $p < 0.001$ and 0.8%, 6 seconds, $p = 0.02$, in 2005–2006 and 2006–2007, respectively) by prior cycling, in both seasons (Table 3, Figure 1).

Changes in Cardiorespiratory Response and Performance over Two Consecutive Seasons

Tables 2 and 3, and Figures 2 and 3, respectively, show the evolution of the cardiorespiratory response to C-R over time, in TID and SE triathletes. Our younger athlete group exhibited an increase ($p < 0.001$) in the mean $\dot{V}O_2$ from season 2005–2006 to 2006–2007 during the running component of the C-R test but not during R. However, cycling $\dot{V}O_{2\max}$ remained stable (no significant differences) between seasons (Table 1). The $\dot{V}_E/\dot{V}O_2$ ratio decreased ($p = 0.02$) in both tests, but $\dot{V}_E/\dot{V}CO_2$ was lower ($p < 0.001$) in the second season only during R (Table 2). Moreover, the mean HR during the R test, but not during the C-R, was lower in season 2006/2007 than in 2005/2006 ($p < 0.001$). On the other hand, SE triathletes presented similar HRs and $\dot{V}O_2$ over both seasons and tests (Table 3, Figure 3).

Although the TID triathletes took less time to complete the time-trial run in season 2006–2007 with respect to 2005/2006, both the C-R (–6.6%, 45 seconds, $p < 0.01$) and the R (–6.4%, 41 seconds, $p = 0.01$) tests (Table 2, Figure 4), 3-km time-trial times remained stable in the SE triathletes over both seasons and the 2 tests conditions (–3.9% in C-R and +0.3% in R) (Table 3, Figure 3). Figure 2 shows that the differences between consecutive seasons in running speed within C-R were concentrated within the first 2,000 m of the run in the younger athletes. Within the R test condition, the differences occurred at the beginning (from 400 to 1,200 m) and near the end (from 2,400 m) of the trial. Analysis of consecutive 400-m laps did not reveal any significant difference between seasons in running speed in the SE triathletes either during C-R or R (Figure 3). In both tests and groups, pacing was similar involving a quicker start in the first lap and a final increase in speed over the last 600 m of the run (Figures 2 and 3). A ‘U-shape’ pacing pattern was more evident in the TID triathletes than in the SE. Individual subject analysis of running performance revealed that only 1 triathlete out of 9 in the TID group did not improve his performance during C-R in the season 2006–2007 over that observed in 2005–2006 (Figure 4, left panels). The SE group displayed a higher intragroup variability and during the C-R test 2 out of 6 triathletes performed worse in 2006–2007. Similarly, during the R test run, half of the SE triathletes improved in the second season of the study, whereas half took more time to complete the time trial than taken previously (Figure 4 right panels).

DISCUSSION

Although almost a decade has elapsed since it was pointed out (29) that longitudinal examination of running performance indices during simulated T2 could prove a useful adjunct to training (6), particularly in the case of young triathletes (25,29), this study appears to be the first to have attempted to verify this. Our results demonstrate that the negative effect of cycling on subsequent running performance, as assessed by the time taken to complete a 3-km time-trial run, was maintained

over the 2 consecutive seasons of the study in both TID and SE triathletes. Nonetheless, running performance did improve in the second season but only in the young (potential Elite) triathletes, whereas SE triathletes presented a plateau in performance. The improvements, rather than being accompanied by changes in the cardiorespiratory response to the exercise task, appear to have been related to changes in pacing during the post cycling run.

In the literature to date, the extent to which the cardiorespiratory response during the running component of a C-R transition was augmented over that of an isolated control run has been inversely related to the athlete’s ability level or experience and training status (5,12,27,28). Our data agree with those of previous studies as an increased ventilatory response was only observed in the TID triathletes. Indirect measures of respiratory efficiency (i.e., $\dot{V}_E/\dot{V}O_2$ and $\dot{V}_E/\dot{V}CO_2$) were also negatively affected by prior cycling (10,11,16,17) in the latter group. Our subjects performed the post cycling run at intensities (see percentage cycling $\dot{V}O_{2\max}$ in Tables 2 and 3) at which respiratory muscle fatigue has been demonstrated to occur (21). It has been suggested, however, that elite triathletes experience smaller changes in their cardiorespiratory mechanics during the post cycling-running component (16). Thus, the increase in the time taken to complete the time-trial run during the C-R test might be related to respiratory muscle fatigue (10,11) in the TID group, only. Surprisingly, time to complete the 3-km time-trial run was impaired by prior cycle in both groups (Tables 2 and 3, Figure 1). In part, this may be because of the novelty of the C-R test conditions to the SE triathletes leading to some “carryover” of the pattern of cycling muscle recruitment to running, which is not seen during the more familiar race-like cycle conditions (8). Because the cardiorespiratory response was similar in both tests, it is possible that the decrease in performance that was exhibited during the C-R test may be then related to the pacing that was adopted within it (15). Hauswirth et al. (15) demonstrated that pacing during the first kilometer at a speed 5% slower than the average pace of a 10-km control run results in improvements of 150 seconds in the overall time. Our results show that both groups started the running component of the C-R test approximately 3% slower than the average speed at which they completed the isolated run. However, although the SE triathletes adopted a “J-shaped” running strategy, the younger triathletes adopted a “U-shaped” strategy (Figures 2 and 3) (1) (with the latter becoming even more evident, after they had had a further year of experience, in 2006–2007). Moreover, the cycling cadence and the RPE values collected at the end of both the C-R and R tests were near maximal in all cases (Tables 2 and 3). Therefore, in agreement with previous data (15), our results suggest that the ability to maintain post cycling initial speed is an important contributor to final performance.

Biomechanical factors and changes in running pattern after cycling have also been proposed to affect running performance in triathlon (14,18,27,28). It is important to note that top-level

triathletes preserve running kinematics after cycling, but some of them suffer from an alteration in leg muscle activity (5,6). In our case, whatever the primary factor(s) determining the negative effect of cycling on subsequent running performance, the analysis of individual data depicted in Figure 1 shows that cycling affected most of the subjects. Therefore, we suggest that the assessment of how effort or speed is distributed over the course of the time-trial run section of the T2 simulation (15,24), and any possible relationship between this and neuromuscular factors, may prove to be an interesting avenue of future research.

In this study, performance under the C-R test conditions (i.e., 30-minute cycling + 3-km time-trial run) only improved over time in the TID triathletes. We observed, however, that improvements in their post cycling-running performance were accompanied by only slight, nonsignificant, changes in cardiorespiratory parameters. Figure 2 shows that the speed attained during the first 2 km during the time-trial run of the C-R test improved in 2006–2007 compared with that in 2005–2006 without a change in the “U-shape” (1) or cadence during previous cycling. Therefore, it may be worth following up the adaptation to the first stages of a running “off the bike” via noninvasive methods such as an assessment of pacing during a field-based trial, particularly in the case of the younger and less experienced athletes. To date, from the 8 triathletes who improved their performance during the C-R test, 5 are representing Spain at international level, which may be preliminary evidence of the sensitivity of our test to detect specific-T2 adaptations.

Assuming an improvement in running performance from the endurance base to the competitive-period of the season, it is plausible that the negative effect of cycling on consecutive running that was exhibited by the SE triathletes would have either decreased or disappeared by June. Because of calendar competitions and the proximity of the 2008 Olympics, the SE triathletes did not follow the same strategy and running training during the season 2006–2007 and 2005–2006. This could be one reason explaining why their running performance was affected by previous cycling differed between seasons. Nonetheless, SE triathletes displayed a higher intragroup variability (Figure 3). The relevance of these 3-km related results to their competitive performances over the 10-km run of the Olympic distance event, or the 5-km run of the Sprint event, for which a bid for inclusion in the Rio 2016 Olympic Games is currently underway, has not been established.

In this study, we defined performance as “time to complete the 3-km time-trial run during either the C-R or the R test” and the assessment of reliability and validation of the test measures against competition performance is required. However, there is a strong evidence of the relation between post cycling-running performance and competition finishing position (24,37,38), and our test is based on previous findings. First of all, we used a cycling section at the intensity corresponding to the Th_{vent} that lasted 30 minutes because this has been shown to produce a modified cardiorespiratory

response from that observed during an isolated control run (10,11,16–18,20). Second, we chose a 3-km time-trial to assess running performance because the said distance has been both implemented before in specific-T2 tests (3,4) and demonstrated to be sufficient to determine the possible effects of neuromuscular fatigue on performance (30). More importantly, the 3-km distance is the most commonly performed (both in competition and for the purposes of TID) by triathletes under-16 years of age in at least 2 of the 8 highest ranked countries in the ITU qualification simulations for the 2012 Olympic Games (www.triathlon.org, accessed on March, 11, 2011). Finally, Laursen et al. (23) have previously obtained coefficients of variation for 1.5- and 5-km time-trial run of 3.3% (intraclass correlation coefficient [ICC] = 0.879) and 2.0% (ICC = 0.953). Similarly, Schabert et al. (33) have reported a coefficient of variation of 2.7% (ICC = 0.90) for self-paced time-trial performance in endurance runners. Our TID group improved their C-R run times (in 8 triathletes out of 9), between the seasons 2005–2006 and 2006–2007, by >6%—more than expected for time trials over similar distances (23,33). Their mean 3-km run speeds during the C-R tests and within race competitions over the course of the study also did not differ significantly. Overall, it appears likely that the results that we obtained for our TID group reflected a “real” change in their specific-T2 adaptation over time, rather than being because of random variation.

The study was also limited by a lack of collection of stride length and frequency data, which could have helped to explain the results. Nevertheless, the monitoring of performance indices during the post cycling running component over time as a means of optimizing individual specific (performance and injury prevention-related) training intervention(s) (5,7), may prove of major importance to young triathletes in particular (25). In this article, we have presented preliminary evidence that our test might be sensitive enough to detect specific-T2 adaptations, especially in potential Elite triathletes. Our article is the first to report longitudinal changes in the cardiorespiratory response and running performance in a field-based C-R transition over the course of 2 consecutive seasons. Although running performance after cycling remained stable in SE Spanish National Squad triathletes, it significantly improved in young triathletes enrolled in the Spanish National Talent Identification Plan. This improvement may have been related to differences in pacing rather than to changes in the cardiorespiratory response per se.

PRACTICAL APPLICATIONS

This work provides further insight into the evolution of performance during cycling-running succession. The existing academic literature and our results suggest that coaches should be aware of the importance of pacing strategy (especially at the beginning of the running component) on final performance. The incorporation of training sessions that are designed to automate pacing strategy into the athlete’s training plan could be a useful means of improving their performance.

Although both the reliability and validity of our test need to be further evaluated, we have presented preliminary evidence that it may be sensitive to specific-T2 adaptation in the youngest triathletes. Once it has been more fully researched, our field-based test, likely more accessible to coaches than laboratory-based tests, may prove to be a useful addition to long-term triathlete development plans.

ACKNOWLEDGMENTS

The authors wish to thank Juan Rodríguez Biehn for his collaboration in this work, which was funded by the Ministerio de Educación y Ciencia and the Consejo Superior de Deportes (projects 04/UPB10/05 and 03/UPB10/07, respectively). V.D. is funded by Universidad Politécnica de Madrid (Action Marie Curie-COFUND, contract UNITE No. 246565). V.E.V. thanks the Fundação para a Ciência e a Tecnologia (the Portuguese Foundation for Science and Technology) for their award of a postdoctoral research fellowship under the Ciência 2008 program. The authors declare no conflict of interests. None of the results of this study constitute endorsement of the product by the authors or the National Strength and Conditioning Association. F.J.C. and A.G.Z. contributed equally to the work.

REFERENCES

1. Abbiss, CR and Laursen, PB. Describing and understanding pacing strategies during athletic competition. *Sports Med* 38: 239–252, 2008.
2. Beaver, WL, Wasserman, K, and Whipp, BJ. A new method for detecting anaerobic threshold by gas exchange. *J Appl Physiol* 60: 2020–2027, 1986.
3. Bernard, T, Verduyssen, F, Grego, F, Hausswirth, C, Lepers, R, Vallier, JM, and Brisswalter, J. Effect of cycling cadence on subsequent 3 km running performance in well trained triathletes. *Br J Sports Med* 37: 154–159, 2003.
4. Bernard, T, Verduyssen, F, Mazure, C, Gorce, P, Hausswirth, C, and Brisswalter, J. Constant versus variable-intensity during cycling: Effects on subsequent running performance. *Eur J Appl Physiol* 99: 103–111, 2007.
5. Bonacci, J, Blanch, P, Chapman, AR, and Vicenzino, B. Altered movement patterns but not muscle recruitment in moderately trained triathletes during running after cycling. *J Sports Sci* 28: 1477–1487, 2010.
6. Bonacci, J, Green, D, Saunders, PU, Blanch, P, Franettovich, M, Chapman, AR, and Vicenzino, B. Change in running kinematics after cycling are related to alterations in running economy in triathletes. *J Sci Med Sport* 13: 460–464, 2010.
7. Bonacci, J, Green, D, Saunders, PU, Franettovich, M, Blanch, P, and Vicenzino, B. Plyometric training as an intervention to correct altered neuromotor control during running after cycling in triathletes: A preliminary randomised controlled trial. *Phys Ther Sport* 12: 15–21, 2011.
8. Bonacci, J, Saunders, PU, Alexander, M, Blanch, P, and Vicenzino, B. Neuromuscular control and running economy is preserved in elite international triathletes after cycling. *Sports Biomech* 10: 49–61, 2011.
9. Borg, G. *Borg's Perceived Exertion and Pain Scales*. Champaign, IL: Human Kinetics, 1998.
10. Boussana, A, Galy, O, Hue, O, Matecki, S, Varray, A, Ramonatxo, M, and Le Gallais, D. The effects of prior cycling and a successive run on respiratory muscle performance in triathletes. *Int J Sports Med* 24: 63–70, 2003.
11. Boussana, A, Matecki, S, Galy, O, Hue, O, Ramonatxo, M, and Le Gallais, D. The effect of exercise modality on respiratory muscle performance in triathletes. *Med Sci Sports Exerc* 33: 2036–2043, 2001.
12. Chapman, AR, Vicenzino, B, Blanch, P, Dowlan, S, and Hodges, PW. Does cycling effect motor coordination of the leg during running in elite triathletes? *J Sci Med Sport* 11: 371–380, 2008.
13. Díaz, V, Benito, PJ, Peinado, AB, Álvarez, M, Martín, C, di Salvo, V, Pigozzi, F, Maffulli, N, and Calderón, FJ. Validation of a new portable metabolic system during an incremental running test. *J Sport Sci Med* 7: 532–536, 2008.
14. Hausswirth, C, Bigard, AX, and Guezennec, CY. Relationships between running mechanics and energy cost of running at the end of a triathlon and a marathon. *Int J Sports Med* 18: 330–339, 1997.
15. Hausswirth, C, Le Meur, Y, Bieuzen, F, Brisswalter, J, and Bernard, T. Pacing strategy during the initial phase of the run in triathlon: Influence on overall performance. *Eur J Appl Physiol* 108: 1115–1123, 2010.
16. Hue, O, Galy, O, Le Gallais, D, and Prefaut, C. Pulmonary responses during the cycle-run succession in elite and competitive triathletes. *Can J Appl Physiol* 26: 559–573, 2001.
17. Hue, O, Le Gallais, D, Boussana, A, Chollet, D, and Prefaut, C. Ventilatory responses during experimental cycle-run transition in triathletes. *Med Sci Sports Exerc* 31:1422–1428, 1999.
18. Hue, O, Le Gallais, D, Chollet, D, Boussana, A, and Prefaut, C. The influence of prior cycling on biomechanical and cardiorespiratory response profiles during running in triathletes. *Eur J Appl Physiol Occup Physiol* 77: 98–105, 1998.
19. Hue, O, Le Gallais, D, Chollet, D, and Prefaut, C. Ventilatory threshold and maximal oxygen uptake in present triathletes. *Can J Appl Physiol* 25: 102–113, 2000.
20. Hue, O, Le Gallais, D, and Prefaut, C. Specific pulmonary responses during the cycle-run succession in triathletes. *Scand J Med Sci Sports* 11: 355–361, 2001.
21. Johnson, BD, Babcock, MA, Suman, OE, and Dempsey, JA. Exercise-induced diaphragmatic fatigue in healthy humans. *J Physiol* 460: 385–405, 1993.
22. Kreider, RB, Boone, T, Thompson, WR, Burkes, S, and Cortes, CW. Cardiovascular and thermal responses of triathlon performance. *Med Sci Sports Exerc* 20: 385–390, 1988.
23. Laursen, PB, Francis, GT, Abbiss, CR, Newton, MJ, and Nosaka, K. Reliability of time-to-exhaustion versus time-trial running tests in runners. *Med Sci Sports Exerc* 39: 1374–1379, 2007.
24. Le Meur, Y, Hausswirth, C, Dorel, S, Bignet, F, Brisswalter, J, and Bernard, T. Influence of gender on pacing adopted by elite triathletes during a competition. *Eur J Appl Physiol* 106: 535–545, 2009.
25. Millet, GP and Bentley, DJ. The physiological responses to running after cycling in elite junior and senior triathletes. *Int J Sports Med* 25: 191–197, 2004.
26. Millet, GP, Dreano, P, and Bentley, DJ. Physiological characteristics of elite short- and long-distance triathletes. *Eur J Appl Physiol* 88: 427–430, 2003.
27. Millet, GP, Millet, GY, and Candau, RB. Duration and seriousness of running mechanics alterations after maximal cycling in triathletes. Influence of the performance level. *J Sports Med Phys Fitness* 41: 147–153, 2001.
28. Millet, GP, Millet, GY, Hofmann, MD, and Candau, RB. Alterations in running economy and mechanics after maximal cycling in triathletes: Influence of performance level. *Int J Sports Med* 21: 127–132, 2000.
29. Millet, GP and Vleck, VE. Physiological and biomechanical adaptations to the cycle to run transition in Olympic triathlon: Review and practical recommendations for training. *Br J Sports Med* 34: 384–390, 2000.

30. Paavolainen, LM, Nummela, AT, and Rusko, HK. Neuromuscular characteristics and muscle power as determinants of 5-km running performance. *Med Sci Sports Exerc* 31: 124–130, 1999.
31. Rabadán, M, Díaz, V, Calderón, FJ, Benito, PJ, Peinado, AB, and Maffulli, N. Physiological determinants of specialty of elite middle- and long-distance runners. *J Sports Sci* 29: 975–982, 2011.
32. Rosdahl, H, Gullstrand, L, Salier-Eriksson, J, Johansson, P, and Schantz, P. Evaluation of the Oxycon Mobile metabolic system against the Douglas bag method. *Eur J Appl Physiol* 109: 159–171, 2010.
33. Schabert, EJ, Hopkins, WG, and Hawley, JA. Reproducibility of self-paced treadmill performance of trained endurance runners. *Int J Sports Med* 19: 48–51, 1998.
34. Smith, DJ. A framework for understanding the training process leading to elite performance. *Sports Med* 33: 1103–1126, 2003.
35. Vleck, V and Alves, F. Triathlon transition tests: Overview and recommendations for future research. *RICYDE: Rev Int Cienc Dep* 7: I–III, 2011.
36. Vleck, V, Santos, S, Bentley, D, and Alves, F. Influence of prior cycling on the OBLA measured during incremental running in triathletes. In: *Annual Congress of the British Association of Sports and Exercise Scientists. Leeds: J Sports Sci*, 23:93–223, 2005.
37. Vleck, VE, Bentley, DJ, Millet, GP, and Burgi, A. Pacing during an elite Olympic distance triathlon: Comparison between male and female competitors. *J Sci Med Sport* 11: 424–432, 2008.
38. Vleck, VE, Burgi, A, and Bentley, DJ. The consequences of swim, cycle, and run performance on overall result in elite Olympic distance triathlon. *Int J Sports Med* 27: 43–48, 2006.