

ORIGINAL ARTICLES

Response of Tidal Volume to Inspiratory Time Ratio During Incremental Exercise

P.J. Benito,^a F.J. Calderón,^a A. García-Zapico,^b J.C. Legido,^c and J.A. Caballero^d

^aFacultad de Ciencias de la Educación Física y del Deporte (INEF), Universidad Politécnica de Madrid, Madrid, Spain.

^bFacultad de Educación, Universidad Complutense de Madrid, Madrid, Spain.

^cEscuela Profesional de Especialistas en Medicina de la Educación Física y del Deporte, Facultad de Medicina, Universidad Complutense de Madrid, Madrid, Spain.

^dFacultad de Ciencias de la Actividad Física y el Deporte, Las Palmas de Gran Canaria, Las Palmas, Spain.

OBJECTIVE: There is some debate about the participation of the Hering-Breuer reflex during exercise in human beings. This study aimed to investigate breathing pattern response during an incremental exercise test with a cycle ergometer. Participation of the Hering-Breuer reflex in the control of breathing was to be indirectly investigated by analyzing the ratio of tidal volume (V_T) to inspiratory time (t_I).

SUBJECTS AND METHODS: The 9 active subjects who participated the study followed an incremental protocol on a cycle ergometer until peak criteria were reached. During exercise, V_T/t_I can be described in 2 phases, separated by activation of the Hering-Breuer reflex (inspiratory off-switch threshold). In phase I, ventilation increases because V_T increases, resulting in a slight decrease in t_I , whereas, in phase II, increased ventilation is due to both an increase in V_T and a decrease in t_I .

RESULTS: The mean (SD) inspiratory off-switch threshold was 84.6% (6.3%) when expressed relative to peak V_T (mean, 3065 [566.8] mL) and 48% (7.2%) relative to the forced vital capacity measured by resting spirometry. The inspiratory off-switch threshold correlated positively ($r=0.93$) with the second ventilatory threshold, or respiratory compensation point.

CONCLUSIONS: The inspiratory off-switch threshold and V_T/t_I are directly related to one another. The inspiratory off-switch threshold was related to the second ventilatory threshold, suggesting that the Hering-Breuer reflex participates in control of the breathing pattern during exercise. Activation of the reflex could contribute by signaling the respiratory centers to change the breathing pattern.

Key words: Breathing pattern. Hering-Breuer reflex. Ventilatory threshold.

Respuesta de la relación volumen corriente-tiempo inspiratorio durante un esfuerzo incremental

OBJETIVO: La participación del reflejo de Hering-Breuer durante el ejercicio en seres humanos es objeto de discusión. El propósito del presente trabajo ha sido estudiar la respuesta del patrón respiratorio durante un esfuerzo incremental en cicloergómetro para comprobar, de forma indirecta, mediante el análisis de la relación volumen corriente-tiempo inspiratorio (V_T/t_I), la participación del reflejo de Hering-Breuer en el control de la respiración.

SUJETOS Y MÉTODOS: Han participado en el estudio 9 sujetos activos que han llevado a cabo un protocolo incremental en cicloergómetro hasta alcanzar criterios máximos. Se ha comprobado que la relación V_T/t_I durante el ejercicio presenta 2 fases con un punto de ruptura, denominado punto de ruptura Hering-Breuer (PHB): fase I, donde el incremento de la ventilación se produce a expensas del aumento del V_I con ligero descenso del t_I , y fase II, durante la cual el incremento ventilatorio se produce tanto por el aumento del V_I como por el descenso del t_I .

RESULTADOS: En el estudio, el PHB se alcanzaba a un valor medio (\pm desviación estándar) del $84,6 \pm 6,3\%$ respecto al máximo valor de V_T ($3.065 \pm 566,8$ ml) y de un $48 \pm 7,2\%$ respecto al valor de la capacidad vital forzada medida en la espirometría de reposo. El PHB se relacionó de forma positiva ($r = 0,93$) con el umbral ventilatorio 2 o umbral de compensación respiratoria.

CONCLUSIONES: Existe relación directa entre el PHB y V_T/t_I . El PHB se relaciona con el umbral ventilatorio 2, de manera que intervendría en el control del patrón ventilatorio durante el ejercicio. La entrada en funcionamiento del reflejo podría contribuir informando a los centros respiratorios para llevar a cabo el cambio de patrón ventilatorio.

Palabras clave: Patrón respiratorio. Reflejo de Hering-Breuer. Umbral ventilatorio.

Correspondence: Dr. P.J. Benito-Peinado.
Facultad de Ciencias de la Educación Física y del Deporte (INEF).
Universidad Politécnica de Madrid.
Martín Fierro, s/n. 28040 Madrid. España.
E-mail: pedroj.benito@upm.es

Manuscript received October 18, 2004. Accepted for publication October 11, 2005.

Introduction

The human breathing pattern has been studied at rest¹ and during both submaximal exercise^{2,4} and maximal exercise.^{2,3,5} All these studies report a response of the ratio of tidal volume (V_T) to inspiratory time (t_I) similar

BENITO PJ ET AL. RESPONSE OF TIDAL VOLUME TO INSPIRATORY TIME RATIO DURING INCREMENTAL EXERCISE

to that described by Clark and von Euler⁶ for animals breathing CO₂-enriched air. These authors described V_T/t_I according to 2 phases. In phase 1, ventilation increases because V_T increases, whereas in phase 2, ventilation increases mainly because t_I decreases.

V_T/t_I can provide an indirect indication of whether the Hering-Breuer reflex has been activated and shows 2 phases well defined by activation of pulmonary receptors, as described by Clark and von Euler.^{6,7} The breathing pattern therefore indicates the transition from breathing in which ventilation increases largely because V_T increases to tachypneic breathing. Such a change in breathing pattern occurs during incremental exercise. In highly-trained cyclists, investigators concur that, above a given intensity, V_T reaches a stable value⁸⁻¹⁰ or increases slightly.¹¹ The breathing rate also participates notably more in the increase in minute ventilation (V_E) above a given intensity of exercise. The contribution of the Hering-Breuer reflex to the control of ventilation during exercise is subject to debate. The breathing pattern of doubly denervated (hilar and carotid-body) animals has been seen to differ to that of normal animals,¹² and recipients of a heart-lung transplant have achieved suitable ventilation during exercise through a large increase of V_T and a decrease in breathing rate,¹³ suggesting that pulmonary receptors are activated during breathing control. These 2 observations, from both animals and humans, indicate that the increase in ventilation during exercise can be partly attributed to information from volume receptors or other types of receptor which communicate with the breathing control centers via the vagal nerve.⁷

The second ventilatory threshold (VT₂) is one of the most widely used variables for spirometric assessment during ergometry. It is defined as the load at which the oxygen uptake is no longer proportional to CO₂ production.¹⁴⁻¹⁶ Given that this variable is sensitive to training,¹¹ we think it should be measured in studies of chronic obstructive pulmonary disease (COPD).

The objective of this study was to analyze the response of breathing pattern and so indirectly determine whether the Hering-Breuer reflex participates in the regulation of breathing and whether activation of this reflex is related to VT₂. According to our initial hypothesis, the change in V_T/t_I during incremental exercise is similar or identical to that described for the first time by Clark and von Euler⁶ during inhalation of CO₂—an experiment that showed vagal activation by means of volume receptors. A relationship between the 2 variables that characterize the breathing pattern could have important applications in COPD.

Subjects and Methods

Subjects

Descriptive statistics of the population are presented in Table 1. Nine amateur male cyclists, triathletes, and physical education students who were regular cyclists were recruited. All gave written informed consent to participate in the study.

The study was conducted in accordance with the ethical principles of the World Medical Association Declaration of Helsinki for investigation with human subjects.¹⁷

Measurement of Gas Exchange

The composition and volume of expired air were measured with a Jaeger Oxycon Pro[®] gas analyzer (Erich Jaeger, Würzburg, Germany). The 2-way digital turbine sensor (Triple V[®], Jaeger) had low dead space and resistance and complied with the guidelines of the American Thoracic Society and the European Community for Steel and Coal.¹⁸ Gas exchange data were collected breath by breath and averaged over 15 seconds. Heart rate was measured with a Hellige Cardiotest EK 53 3-lead electrocardiograph (Hellige, Freiburg, Germany).

Study Protocol

On arrival at the laboratory, each subject underwent a medical examination, which consisted of a medical and sporting history, basal electrocardiogram, and spirometry testing. The tests were done on a Jaeger ER800[®] variable resistance cycle ergometer with electromagnetic braking. Loads covered by the ergometer ranged from 25 W to 1000 W with minimum increments of 1 W/s. The exercise protocol consisted of 1 minute at full rest, 3 minutes warm-up at 50 W, and progressive testing with increments of 5 W every 12 seconds. At the end of the exercise program, the subjects had 2 minutes of active recovery at 50 W (70 rpm), and 3 minutes of complete rest on the bicycle. The pedal rate was set at between 70 rpm and 90 rpm. All tests were done under similar atmospheric conditions (temperature range, 21°C to 24°C; relative humidity, 45% to 55%; and atmospheric pressure, 700 mm Hg to 715 mm Hg). Values were expressed in standard temperature and pressure dry conditions.

Analysis of Anaerobic Threshold

Data on gas exchange were analyzed by visual inspection as proposed by Wassermann et al¹⁹ in 1973. In accordance with this method, V_E/CO_2 production, V_E /oxygen uptake, partial pressure of CO₂, and partial pressure of oxygen were measured at different loads to determine the anaerobic threshold using standard criteria. A combination of different methods has been shown to be the most reliable to determine ventilatory thresholds.¹⁵

For each subject, the values of V_T from the exercise test were plotted against t_I in order to visualize the 2 phases of the breathing pattern equivalent to those reported by Clark and von Euler during inhalation of CO₂.⁶ In phase 1, increases in ventilation are due to increases in V_T with a slight decrease in t_I . In phase 2, the ventilatory increase occurs because of both increases in V_T and decreases in t_I , with an inverse relationship. This variation in ventilatory response is reached when V_T is about 2 times the resting value.

In the Clark and von Euler breathing pattern, the phase transition occurs suddenly and is known as the inspiratory off-switch threshold. Visual examination reveals the point at which V_T/t_I shifts to the left on the plot, that is, the point where t_I decreases significantly with a small increase in V_T . Two independent observers, blinded to the identity of the subject, determined VT₂ and the inspiratory off-switch threshold at different times.

BENITO PJ ET AL. RESPONSE OF TIDAL VOLUME TO INSPIRATORY TIME RATIO DURING INCREMENTAL EXERCISE




Figure 1. Left panel: ratio of oxygen uptake (VO_2) on reaching the inspiratory off-switch threshold (IOF) to VO_2 on reaching the second ventilatory threshold (VT_2). Right panel: Bland and Altman procedure²⁰ (1986) for analysis of the reliability of the procedure for identification of VT_2 . Diff indicates difference.

Statistical Analysis

Mean (SD) values were calculated for all variables. After testing for a normal distribution, the Student *t* test was applied for paired variables. A Pearson correlation study was used to test whether there was a positive relationship between the anaerobic threshold and the inspiratory off-switch threshold. The Bland and Altman procedure²⁰ was used to investigate the

TABLE 1
Characteristics of the Subjects Included in the Study*

	Mean (SD)
Age, years	24.1 (3)
Weight, kg	70.5 (6.5)
Height, cm	176 (6.6)
FVC, L	5.4 (0.9)
FEV ₁ , L	4.6 (0.6)
VO ₂ , mL×kg×min ⁻¹	57.1 (12.1)
Maximum load, W	336 (8)

*FVC indicates forced vital capacity; FEV₁, forced expiratory volume in 1 second; VO₂, peak oxygen uptake adjusted to body weight. Maximum load is defined as the mean peak load reached during the tests.

TABLE 2
Results of the Analysis of the Ratio of Tidal Volume (V_T) to Inspiratory Time (t_I) at Peak Exercise and Analysis of Comparison of Means*

	VO ₂ IOF, mL×min ⁻¹	VO ₂ VT ₂ , mL×min ⁻¹	V _T IOF, mL	Peak V _T , mL
Mean	3103	3375†	2576	3065
SD	751	637	441	566
	V _T OIF/ Peak V _T	V _T OIF/ FVC	Peak V _T / FVC	Peak V _T / FEV ₁
Mean	84.6%	48.0%	56.8%	66.1%
SD	6.3%	7.2%	7.6%	8.1%

*VO₂ IOF indicates oxygen uptake on reaching the inspiratory off-switch threshold; VO₂ VT₂, oxygen uptake on reaching the second ventilatory threshold; V_T IOF, tidal volume on reaching the inspiratory off-switch threshold; FVC, forced vital capacity; FEV₁, forced expiratory volume in 1 second. †Significant differences VO₂ IOF versus VO₂ VT₂: *t* (8)=2.559; *P*=.034.

reliability of the V_T/t_I plots for determining VT_2 . The results of statistical tests were significant when *P* was less than .05. For statistical analysis, the SPSS program, version 11.5 (SPSS Worldwide Headquarters, Chicago, Illinois, USA) was used.

Results

Table 1 shows the anthropometric data, resting spirometric variables, and peak oxygen uptake in the study population. The values of peak oxygen uptake (mean 57.1 [12.4] mL×kg×min⁻¹) were greater than those reported for the sedentary population. A maximum load of 336 W was reached. Table 2 shows the corresponding values at the moment when the breathing pattern changed phase. The 2 phases described in the previous section could be discerned in the exercise tests of all volunteers.

The inspiratory off-switch threshold occurred when t_I was 0.96 (0.11) seconds and V_T was 2576 (441) mL, equivalent to 84.6% (6.3%) of peak V_T (3065 [566] mL) at maximum exercise intensity, and 48% (7.2%) of peak forced vital capacity measured at rest by spirometry (5416 [823] mL). Peak V_T during the test was 56.8% (7.6%) of the forced vital capacity.

The inspiratory off-switch threshold correlated positively ($r=0.93$; $P=.02$) with VT_2 (Figure 1). According to the Bland and Altman test, only 1 subject had a discrepancy between the 2 measurements. Table 2 shows mean oxygen uptake when the inspiratory off-switch threshold was reached. The threshold occurred at a mean oxygen uptake approximately 300 mL less than that when VT_2 occurred.

Discussion

According to the results of this study, the pattern of V_T/t_I during incremental exercise is similar to that observed by Clark and von Euler in animals forced to

BENITO PJ ET AL. RESPONSE OF TIDAL VOLUME TO INSPIRATORY TIME RATIO DURING INCREMENTAL EXERCISE




Figure 2. Plots of the ratio between tidal volume and inspiratory time for the subjects who participated in the study. The lines correspond to regression fits to determine the inspiratory off-switch threshold. Individual scales are adjusted to illustrate the threshold more clearly.

breathe CO₂.⁶ The response of the ratio shows 2 well differentiated phases (Figure 2). In phase 1, V_E increases mainly because V_T increases and t_I decreases. In contrast, in phase 2, hyperventilation occurs as a result of a greater reduction in t_I with a slight increase in V_T. The moment when the phase change occurs is known as the inspiratory off-switch threshold, which, in this study, is related to VT₂. There is consensus that, above a certain point, V_T reaches a stable value^{9,10,21} or increases slightly,¹¹ according to the subject's level of fitness. Some authors have shown that high intensity strength training can increase V_T in patients with COPD.²² No change in the breathing pattern at rest was observed by Kay et al,²³ however; nor has change been seen under anesthetic,¹⁰ although changes have been reported in recipients of a heart-lung transplant.^{13,24} The findings of Kay et al²³ may differ from ours because the subjects in their study followed an exercise protocol with a limited load of 50 W to 200 W, whereas we used a incremental protocol that reached the maximum load.

Given that our study was with human subjects, it was not possible to conclusively demonstrate that vagal reflexes were participating in ventilatory control during exercise because we could not control for many of the variables that might influence the behavior of V_T/t_I. The physiological mechanisms that may regulate breathing during exercise can be classed as feedforward, feedback, and short-term potentiation,²⁵ although such mechanisms would not completely explain the response of V_E. Feedback could arise from structures in the central nervous system structures (central feedback) or receptors located in the respiratory tract (parenchyma and thorax), in the locomotor muscles, or in the cardiovascular system (peripheral chemoreceptors), that is, peripheral feedback. Therefore, we could not control information from the implicated group III muscle nerve endings, peripheral chemoreceptors, or receptors sensitive to certain molecules and ions (potassium, protons).

Despite these limitations, the results of the study show that V_T/t_I followed a biphasic pattern during

BENITO PJ ET AL. RESPONSE OF TIDAL VOLUME TO INSPIRATORY TIME RATIO DURING INCREMENTAL EXERCISE

incremental exercise, the inspiratory off-switch threshold being reached when V_T was 84.6% (6.3%) of the peak value. Such a pattern could be explained by activation of the Hering-Breuer reflex. This reflex mechanism has been shown to participate in control of breathing in animal models.²⁶ Activation of the Hering-Breuer reflex is suggested by the fact that the inverse relationship between V_T and t_i is affected in a variety of experimental conditions (vagotomy, hilar denervation, hypercapnia, β -receptor blockade, and mouth pressure) both in animals and human beings.

Vagotomy causes an increase in V_E because V_T increases with a minimum decrease in t_i , such that V_T/t_i would be in phase 1.^{13,27} In an interesting study that aimed to demonstrate the effects of double denervation (hilar denervation and carotid chemoreceptor denervation), lung afferents via the hilar nerves influenced the breathing pattern of ponies at rest and during exercise.¹² The effect was also associated with attenuation of lung volume feedback.

There is vagal participation in control of breathing when hypercapnia occurs during exercise in both animals²⁸ and humans.⁹ Joyner et al²⁹ showed that blockade of β_1 - and β_2 -receptors led to larger decreases in V_T than blockade of β_1 -receptors only.

Studies done using the simplest technique, that is, mouth pressure at 0.1 seconds after onset of inspiration^{5,30,31} to determine V_T/t_i , showed that increases in ventilation during easy exercise occurred due to increases in V_T .

In view of the findings of these studies, our results point to the importance of information from airway receptors and receptors in the lung parenchyma. The change in breathing pattern after passing the inspiratory off-switch threshold could be due to feedback from volume receptors or another type of receptor. Such receptors would be partly responsible for the decrease in t_i when V_T increases. The role of the Hering-Breuer reflex in regulation of breathing at rest has been extensively debated,²⁶ as the receptors not only detect changes in volume, but also variations in the concentrations of certain molecules. Other physiological mechanisms, such as variations in CO_2 concentration and increase in the values of certain other variables may therefore explain the results obtained for the inspiratory off-switch threshold.

Slight changes in partial pressure of CO_2 are understood to affect the stimulation of receptors implicated in the Hering-Breuer reflex.³² Stimulated receptors could trigger a strong activation of the bulbar centers by way of vagal inputs during breathing. In our study, given that the inspiratory off-switch threshold is related to VT_2 , above which the partial pressure of exhaled CO_2 increases, variations in partial pressure of CO_2 could have occurred which would explain stimulation of pulmonary receptors sensitive to this gas.

At high intensities, the concentration of plasma protons increases (metabolic acidosis) and potassium concentration varies, sometimes reaching values as high

as 7 mEq/L. This would partly explain hyperventilation during exercise, whether by stimulation of peripheral or central chemoreceptors.^{33,34} We did not measure variables to determine the acid-base status, and so there is no way of knowing whether the concentration of potassium and protons in blood could have led to the change in breathing pattern. Nevertheless, the partial pressure of CO_2 changes after exceeding the inspiratory off-switch threshold, such that slight variations in this variable could explain stimulation of volume receptors or other types of CO_2 receptors.³⁵

Analysis of VT_2 can also be used to determine V_T/t_i , although caution should be exercised, as the inspiratory off-switch threshold is exceeded at a value of approximately 300 mL lower. The inspiratory off-switch threshold very likely forms part of the process of anticipation of increased ventilation. Our measurements of the ratio of V_T to forced vital capacity agree with those of other authors who have measured this variable and who also found statistically significant differences between those with and without COPD.³⁶ This variable has also proved to be a good indicator of the improvement in ventilation with training.²² Therefore, the measurement of the inspiratory off-switch threshold could be a useful method for following the progress of patients with COPD.

In conclusion, the behavior of V_T/t_i during incremental exercise can be divided into 2 phases. The first of these phases shows an increase in ventilation because V_T increases and, to a lesser extent, breathing rate increases (that is, t_i decreases). The second phase shows increased breathing rate, so limiting further increases in V_T . The phase transition occurs at the inspiratory off-switch threshold, which is related to VT_2 . While aware of the limitations of the study, we suggest that the inspiratory off-switch threshold may arise because of stimulation of volume receptors, such that information is sent to the regulatory centers to increase ventilation at a given extent of pulmonary distension. Measurement of the threshold could be useful in patients with COPD.

REFERENCES

1. von Euler C. On the central pattern generator for the basic breathing rhythmicity. *J Appl Physiol.* 1983;55:1647-59.
2. Askanazi J, Milic-Emili J, Broell JR, Hyman AI, Kinney JM. Influence of exercise and CO_2 on breathing pattern of normal man. *J Appl Physiol.* 1979;47:192-6.
3. Clark JM, Hagerman FC, Gelfand R. Breathing patterns during submaximal and maximal exercise in elite oarsmen. *J Appl Physiol.* 1983;55:440-6.
4. Ellingsen I, Sydnes G, Hauge A, Nicolaysen G. Effects of exercise and CO_2 inhalation on the breathing pattern in man. *Acta Physiol Scand.* 1988;134:161-73.
5. Lind F, Hesser CM. Breathing pattern and lung volumes during exercise. *Acta Physiol Scand.* 1984;120:123-9.
6. Clark FJ, von Euler C. On the regulation of depth and rate of breathing. *J Physiol.* 1972;222:267-95.
7. Dempsey JA, Adams L, Ainsworth D, Fregosi RF, Gallagher CG, Guz A, et al. Airway, lung and respiratory muscle function during exercise. In: Chenuck NS, Widdicombe JG, editors. *Handbook of physiology exercise: regulation and integration of multiple systems.* Bethesda: American Physiological Society; 1986. p. 69-91.

BENITO PJ ET AL. RESPONSE OF TIDAL VOLUME TO INSPIRATORY TIME RATIO DURING INCREMENTAL EXERCISE

8. Gallagher CG, Brown E, Younes M. Breathing pattern during maximal exercise and during submaximal exercise with hypercapnia. *J Appl Physiol.* 1987;63:238-44.
9. Martin BJ, Weil JV. CO₂ and exercise tidal volume. *J Appl Physiol.* 1979;46:322-5.
10. Polacheck J, Strong R, Arens J, Davies C, Metcalf I, Younes M. Phasic vagal influence on inspiratory motor output in anesthetized human subjects. *J Appl Physiol.* 1980;49:609-19.
11. Lucía A, Carvajal A, Calderón FJ, Alfonso A, Chicharro JL. Breathing pattern in highly competitive cyclists during incremental exercise. *Eur J Appl Physiol Occup Physiol.* 1999;79:512-21.
12. Flynn C, Forster HV, Pan LG, Bisgard GE. Role of hilar nerve afferents in hyperpnea of exercise. *J Appl Physiol.* 1985;59:798-806.
13. Scirba FC, Owens GR, Sanders MH, Griffith BP, Hardesty RL, Paradis IL, et al. Evidence of an altered pattern of breathing during exercise in recipients of heart-lung transplants. *N Engl J Med.* 1988;319:1186-92.
14. Beaver WL, Wasserman K, Whipp BJ. A new method for detecting anaerobic threshold by gas exchange. *J Appl Physiol.* 1986;60:2020-7.
15. Gaskill SE, Ruby BC, Walker AJ, Sánchez OA, Serfass RC, Leon AS. Validity and reliability of combining three methods to determine ventilatory threshold. *Med Sci Sports Exerc.* 2001;33:1841-8.
16. Wasserman K, Hansen JE, Sue D Y, Whipp BJ, Casaburi R. Principles of exercise testing and interpretation. 2nd ed. Philadelphia: Lea & Febiger; 1994.
17. WMA. Declaración de Helsinki para la investigación con seres humanos. 2000. Available from: <http://www.wma.net/s/policy/b3.htm>
18. Quanjer GJ, Tammeling JE, Cotes LM, Fabbri H, Matthys OF, Pedersen R et al. Lung volume and forced ventilatory flows. Report working party standardization of lung function tests, European Community for Steel and Coal. *Eur Respir J Suppl.* 1993;16:5-40.
19. Wasserman K, Whipp BJ, Koyle SN, Beaver WL. Anaerobic threshold and respiratory gas exchange during exercise. *J Appl Physiol.* 1973;35:236-43.
20. Bland JM, Altman DJ. Regression analysis. *Lancet.* 1986;19:908-9.
21. Gallagher CG. Exercise and chronic obstructive pulmonary disease. *Med Clin North Am.* 1990;74:619-41.
22. Ruiz de Ona Lacasta JM, García de Pedro J, Puente Maestu L, Llorente Íñigo D, Celdrán Gil J, Cubillo Marcos JM. Effects of muscle training on breathing pattern in patients with severe chronic obstructive pulmonary disease. *Arch Bronconeumol.* 2004;40:20-3.
23. Kay JD, Petersen ES, Vejby-Christensen H. Breath-by-breath pattern in man during steady-state bicycle exercise. *J Physiol.* 1975;244:52P-3P.
24. Scirba FC, Owens GR, Sanders MH, Costantino JP, Paradis IL, Griffith BP. The effect of obliterative bronchiolitis on breathing pattern during exercise in recipients of heart-lung transplants. *Am Rev Respir Dis.* 1991;144:131-5.
25. Eldridge FL, Waldrop TG. Neural control of breathing during exercise. In: Whipp BJ, Wasserman K, editors. Exercise: pulmonary physiology and pathophysiology. New York: Marcel Dekker, Inc.; 1991. p. 309-70.
26. Coleridge HM, Coleridge JCG. Reflexes evoked from tracheobronchial tree and lungs. In: Fishman AP, editor. Handbook of physiology. Bethesda: American Physiological Society; 1986. p. 407-13.
27. Clifford PS, Litzow JT, von Colditz JH, Coon RL. Effect of chronic pulmonary denervation on ventilatory responses to exercise. *J Appl Physiol.* 1986;61:603-10.
28. Ainsworth DM, Smith CA, Johnson BD, Eicker SW, Henderson KS, Dempsey JA. Vagal modulation of respiratory muscle activity in awake dogs during exercise and hypercapnia. *J Appl Physiol.* 1992;72:1362-7.
29. Joyner MJ, Jilka SM, Taylor JA, Kalis JK, Nittolo J, Hicks RW, et al. Beta-blockade reduces tidal volume during heavy exercise in trained and untrained men. *J Appl Physiol.* 1987;62:1819-25.
30. Milic-Emili J, Zin WA. Relationship between neuromuscular respiratory drive and ventilatory output. In: Fishman AP, Macklem PT, Mead J, Geiger SR, editors. Handbook of physiology. The respiratory system. Bethesda: American Physiological Society; 1986. p. 631-46.
31. Whitelaw WA, Derenne JP, Milic-Emili J. Occlusion pressure as a measure of respiratory center output in conscious man. *Respir Physiol.* 1975;23:181-99.
32. Matsuoka T, Mortola JP. Effects of hypoxia and hypercapnia on the Hering-Breuer reflex of the conscious newborn rat. *J Appl Physiol.* 1995;78:5-11.
33. McLoughlin P, Popham P, Linton RA, Bruce RC, Band DM. Exercise-induced changes in plasma potassium and the ventilatory threshold in man. *J Physiol.* 1994;479:139-47.
34. Paterson DJ. Potassium and breathing in exercise. *Sports Med.* 1997;23:149-63.
35. Kaufman MP, Forster HV. Reflexes controlling circulatory. Ventilatory and airway responses to exercise. In: Rowell LB, Mitchell JH, editors. Handbook of physiology. Exercise: regulation and integration of multiple systems. New York: Oxford University Press; 1996. p. 333-80.
36. González-García M, Barrero M, Maldonado D. Exercise limitation in patients with chronic obstructive pulmonary disease at the altitude of Bogota (2,640 m). Breathing pattern and arterial gases at rest and peak exercise. *Arch Bronconeumol.* 2004;40:54-61.